

Conference Highlights ¹

COSMIC FLOWS: TOWARDS AN UNDERSTANDING OF THE LARGE-SCALE STRUCTURE IN THE UNIVERSE

MICHAEL J. HUDSON

Department of Physics & Astronomy, University of Victoria, P.O. Box 3055, Victoria, B.C. V8W 3P6, Canada. E-mail: hudson@uvastro.phys.uvic.ca

This workshop was the first meeting devoted exclusively to “Cosmic Flows” since the conference held at the Institut d’Astrophysique de Paris exactly 6 years earlier. The major issues addressed in Victoria were summarized by Strauss in his introductory review:

1. On the largest scales, have we identified a volume which is at rest with respect to the Cosmic Microwave Background (CMB)?
2. What can be learned from the “coldness” of the flow field on small scales?
3. Can we agree on a value of $\beta \equiv \Omega^{0.6}/b$? If not, is it because the relationship between mass and light (“biasing”) is not a simple linear function? How can peculiar velocity studies be used to constrain biasing?
4. What are the next generation of peculiar velocity surveys?

1. SEEKING CONVERGENCE TO THE CMB FRAME

One of the long-standing goals of observational cosmology has been to identify a fair sample of the Universe. Within the context of cosmic flows, this amounts to identifying the depth at which the local Universe is at rest with respect to the CMB, and thus contains all of the major attractors responsible for the 600 km s^{-1} motion of the Local Group. In practice, this can be done by selecting a sample of galaxies for which distances are known and determining their net bulk peculiar velocity in the CMB frame.

Coming into this workshop, the situation to a depth of $\lesssim 50h^{-1} \text{ Mpc}$ was fairly clear: existing peculiar velocity catalogs indicated that the CMB-frame bulk motion of this volume is $\sim 300 \pm 100 \text{ km s}^{-1}$. A tidal analysis (Dekel)² indicates that most of this motion is due to masses beyond $\lesssim 50h^{-1} \text{ Mpc}$. However, newly-presented observational results on these scales favor the lower end of this range, and thus more local sources. Tonry & Dressler obtained precise distances to nearby ($\lesssim 30 h^{-1} \text{ Mpc}$) galaxies with the Surface Brightness Fluctuation (SBF) technique. They modeled the flow field with Virgo and a Great Attractor (GA) and found a GA infall velocity of $289 \pm 137 \text{ km s}^{-1}$ at the position of the LG. (This is about half of that originally claimed by the 7 Samurai.) Furthermore, Tonry & Dressler claim only a small ($\lesssim 200 \text{ km s}^{-1}$) residual motion arising from sources beyond their volume. The ENEAR project,

a Fundamental Plane (FP)-based ‘son-of 7-Samurai’ survey, finds somewhat intermediate results: a bulk flow of $343 \pm 53 \text{ km s}^{-1}$ at a depth of $\sim 50h^{-1} \text{ Mpc}$ (Wegner). The ‘Shellflow’ project (Courteau) using the Tully-Fisher (TF) distance indicator, found that galaxies in an all-sky shell centered on $55 h^{-1} \text{ Mpc}$ have a low bulk motion of $80 \pm 150 \text{ km s}^{-1}$. Taken together these new results suggest that most of the mass responsible for the motion of the LG lies within $60 h^{-1} \text{ Mpc}$, and can presumably be identified with the GA. Woudt showed results from galaxy redshift surveys in the Zone of Avoidance which indicate an excess of galaxies in the GA region centered on Abell 3627 at $l = 325^\circ$, $b = -7^\circ$. A3627 appears to have the richness of Coma, and its position coincides with the peak of the GA as inferred from POTENT reconstructions.

The observational situation on larger ($> 60h^{-1} \text{ Mpc}$) scales remains complicated. At the ’93 Paris meeting, the controversial results of Lauer & Postman (LP) were dissected. In Victoria, a number of recent results probing similar scales were presented; these were based mainly on sparse samples of galaxy clusters. The FP-based SMAC sample (Smith) and the TF-based LP10k sample (Willick) both find a large bulk flow, $\sim 600 \pm 200 \text{ km s}^{-1}$ (but in a direction different from LP), whereas the TF-based SCI+II (Giovannelli & Dale) sample yields a very low bulk motion ($\sim 100 \pm 100 \text{ km s}^{-1}$). The nearby SNIa (Riess) give an intermediate result $\sim 270 \text{ km s}^{-1}$, in good directional agreement with SMAC/LP10k. The EFAR survey (Colless) finds low peculiar motions, but is restricted to two regions in the sky which do not sample well the directions of the SMAC/LP10k flows. One might wonder if these different groups measure the same peculiar velocities for common clusters. The answer is a qualified yes: formally – given the large errors – the χ^2 is acceptable, but there may be a trend for some surveys have systematically lower peculiar velocities than others (Lucey). It is important to note that what is quoted is the bulk flow of a given sparse sample and not that of a volume. I argued that given the sampling and the presence of internal flows — which act like an extra noise term — all of these surveys (with the exception of LP) are consistent with each other and with a mean large-scale bulk flow of $\sim 300 - 400 \text{ km s}^{-1}$. This is somewhat larger than the bulk motions obtained on smaller scales, but the difference may not be statistically significant. Ultimately, larger and denser samples will be needed to determine the contribution from the largest scales.

Peculiar velocity data can be used to place direct constraints on the mass power spectrum on large scales. Ze-

¹ Conference was held in Victoria, BC, Canada in July 1999. Proceedings will be edited by Stéphane Courteau, Michael Strauss & Jeffrey Willick and published in the *ASP Conference Series*.

²Throughout this summary only the individual who presented results in Victoria is referenced.

havi showed that peculiar velocity data sets gave the constraint $\Omega^{0.6}\sigma_8 \sim 0.8$, somewhat higher than the cluster normalization.

In order to probe the mass power spectrum on small scales, a statistic such as the r.m.s. velocity dispersion of galaxies around the large-scale flow can be used. Baker showed that the 1D single-object weighted velocity dispersion of Las Campanas redshift survey (LCRS) galaxies is $\sim 125 \text{ km s}^{-1}$, which favors $\Omega^{0.5}/b \sim 0.3\text{--}0.6$. This dispersion is similar to the velocity scatter of SBF galaxies around the best fitting flow models (180-200 km s^{-1} , Blakeslee; Tonry & Dressler). Recent theoretical developments on the pairwise infall velocity (Juskiewicz) yield constraints on Ω and σ_8 which are consistent with those obtained from the abundance of rich clusters. It remains to be determined, however, how much the observed small-scale velocity dispersion depends on the biasing of galaxies with respect to dark matter.

2. MASS, LIGHT AND “BIASING”

On large ($> 10h^{-1} \text{ Mpc}$) scales, there appears to be good agreement between mass and light. Dekel showed new POTENT and Weiner Filter reconstructions of the local mass-density field — based on the SBF distances of Tonry et al. — which were in good qualitative agreement with Tully’s galaxy maps of the nearby Universe. When fluctuations in the mass and galaxy densities are much less than the mean density, as they are on large scales, it is reasonable to parameterize the ratio of fluctuations by a single “biasing” parameter, b . A comparison of mass and light galaxies then yields a value of $\beta \equiv \Omega^{0.6}/b$.

There was considerable discussion of β estimates derived from different methods of comparing mass and light. Comparisons of predicted and observed peculiar velocities consistently yield low values $\beta_{\text{IRAS}} \sim 0.45$ (e.g. Blakeslee; Riess). On the other hand, the comparison of the galaxy density with mass density reconstructed from POTENT yields $\beta_{\text{IRAS}} \sim 0.9^3$. Both methods have been extensively tested with N-body simulations and mock catalogs (e.g. Kolatt’s POTENT error analysis). It is possible that this discrepancy is due to the different scales probed: velocity-velocity comparisons are performed on small smoothing scales, whereas the density comparison is smoothed on larger scales. If this is coupled with a scale-dependent biasing factor, as indicated by some semi-analytic models of galaxy formation, then it might account for the different results.

Indeed, on small (few $h^{-1} \text{ Mpc}$) scales, the true relation between mass and light is complicated (Dekel); it is likely to be non-linear, a function of scale, galaxy type, luminosity and surface brightness and may depend on parameters other than the local mass density. While the clustering of

dark matter haloes is now quite well understood through N-body simulations (Frenk; Lake; Klypin) and analytic work (Sheth), the star formation history of a given halo is more uncertain. Pioneering attempts to quantify and explore non-trivial biasing using LCRS and IRAS PSC redshift survey (PSCz) data were presented by Blanton and Narayanan, respectively. Shaya argued that the best fit flow model for the nearby ($\lesssim 30h^{-1} \text{ Mpc}$) Universe was one in which early-type galaxies in dense cores were assigned a mass-to-light ratio of 750 whereas spirals in the field have $M/L \sim 150$. Saunders commented that, in the PSCz catalog, the Hercules and Shapley superclusters are of comparable overdensity, whereas if one counts rich clusters, the latter is far more overdense. Indeed, preliminary POTENT reconstructions of SMAC+SN Ia data indicate a large mass near Shapley (Dekel). It is clear, however, that much more work needs to be done in order to understand the sensitivity of various statistics to simple modifications of the linear biasing scheme.

3. THE FUTURE

Can we improve on current data sets? The 2 Micron All-Sky Survey will obtain redshifts for $\sim 10^5$ galaxies (Huchra) and should provide superlative all-sky galaxy maps for comparison with peculiar velocity surveys. This effort will be complemented by the 6dF survey in Australia which will also obtain velocity dispersions and thus peculiar velocities for all E/S0 galaxies in the South (Mamon). Furthermore, several of the TF/FP teams present in Victoria have plans to extend their cluster or field-based studies.

Other methods also show promise. At present, SN Ia searches are turning up ~ 20 local ($cz < 30000 \text{ km s}^{-1}$) per year, but this should increase with new searches underway (Aldering). The advent of 8m-class telescopes with good seeing will allow the SBF technique to be pushed to $\sim 6000 \text{ km s}^{-1}$, allowing the definitive study of the local flow field. Finally there remains the possibility of using the X-ray emission from clusters as a distance indicator at low redshift, and the kinematic Sunyaev-Zel’dovich effect to probe cluster peculiar velocities at high redshift. When these new data are compared with sophisticated mock catalogs based on semi-analytic galaxy formation models (Dekel), important insights will be gained regarding the relationship between dark and luminous matter.

Overall, the most rewarding aspects of this workshop were the thoughtful questions and very frank responses, both to individual presentations and in the panel-led discussions. These discussions raised more questions than they answered, as they should in any fruitful field. Stéphane Courteau is to be congratulated for organizing a very successful and stimulating meeting.

³The power spectrum analysis of Zehavi yields even higher values: corresponding to $\beta_{\text{IRAS}} \sim 1$, if $b_{\text{IRAS}}\sigma_8 \sim 0.75$.